

CONF-840255--17

Los Alamos National Laboratory is operated by the University of California for the United States Department of Energy under contract W-7405-ENG-38

LA-UR--84-524

DE84 007485

TITLE: THE SUSTAINMENT DYNAMO REEXAMINED: NONLOCAL ELECTRICAL CONDUCTIVITY
OF PLASMA IN A STOCHASTIC MAGNETIC FIELD

AUTHOR(S): Abram R. Jacobson and Ronald W. Moses

SUBMITTED TO: US-Japan Symposium on Compact Toroid Research
Princeton Plasma Physics Laboratory
February 20-23, 1984

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

By acceptance of this article, the publisher recognizes that the U.S. Government retains a nonexclusive, royalty-free license to publish or reproduce the published form of this contribution, or to allow others to do so, for U.S. Government purposes.

The Los Alamos National Laboratory requests that the publisher identify this article as work performed under the auspices of the U.S. Department of Energy

Los Alamos Los Alamos National Laboratory
Los Alamos, New Mexico 87545

THE SUSTAINMENT DYNAMO REEXAMINED: NONLOCAL ELECTRICAL CONDUCTIVITY OF PLASMA IN A STOCHASTIC MAGNETIC FIELD

Abram R. Jacobson and Ronald W. Moses
Los Alamos National Laboratory
Los Alamos, New Mexico 87545 USA

The "plasma dynamo" is both an intriguing and a practical concept. The intrigue derives from attempting to explain naturally occurring¹ and man-made^{2,3} plasmas whose strong field-aligned currents j_{\parallel} apparently disobey the most naive Ohm's law $j_{\parallel} = \sigma_{\parallel} E_{\parallel}$. The practical importance derives from the dynamo's role both in formation and in sustainment of reversed-field pinch (RFP)² and Spheromak³ fusion plasmas. We will examine certain features of the documented quasi-steady discharges² on ZT-40M, an RFP in apparent need⁴ of a sustainment dynamo. We will show that the tail electrons (which carry j_{\parallel}) are probably wandering (along stochastic \vec{B} -field lines) over much of the minor radius in one mean-free-path. This will void any local Ohm's law, whether naive ($j_{\parallel} = \sigma_{\parallel} E_{\parallel}$) or containing additional terms (such as the $\langle \vec{v} \times \vec{B} \rangle_{\parallel}$ of nonlinear dynamo theory). Instead, we will show that observed quasi-steady RFP discharges in ZT-40M are explainable in simple terms ($f = ma$) of electron-momentum diffusion in a stochastic field, using a stochasticity inferred from observed τ_{Ee} . We will then present results of a formal model of this momentum diffusion. The model predicts the key observed anomalies of sustained RFP behavior (excess loop resistance; slower-than-classical current decay) in terms of electron dynamics in a stochastic magnetic field. Absent from our model are the usual turbulent-dynamo concepts: magnetic-helicity conservation, mode-mode interactions, relaxation, wavenumber cascades, etc.

Quasi-steady discharges that defy a naive Ohm's law⁴ have been reported² on ZT-40M. Their parameter regime is low density ($n \leq 2 \times 10^{19} \text{m}^{-3}$), high temperature ($T_e \leq 150 \text{ eV}$), and electron heat-loss time $\tau_{Ee} \approx 10^{-4} \text{ s}$. At moderate pinch parameter ($\theta \leq 1.5$) these RFP discharges show very little poloidal variation of the reversed toroidal field $[B_{\phi}(a)]$ apart from the factor $1/R$: $[\Delta B_z(a)/B_z(a)]_{\text{rms}} \leq 0.1$ and $[\Delta B_z(a)/B_{\theta}(a)]_{\text{rms}} \leq 0.01$. This observed laminarity does not appear to be consistent with the sustainment dynamo's properties seen in MHD calculations by Sykes and Wesson⁵ and by Aydemir and Barnes,⁶ both of which calculations predict^{7,8} such large-scale poloidal asymmetry that $B_z(a)$ is not even everywhere reversed, i.e., $[\Delta B_z(a)/B_z(a)] \sim 1$.

Rechester and Rosenbluth⁹ showed that a typical Tokamak can be driven stochastic (i.e., islands overlap everywhere) with $(B_{\text{r}}^{\text{local}}/B_0)_{\text{rms}} \geq 10^{-5}$ if a wavenumber spectrum populated out to $k_{\perp} \rho_{ci} \approx 1$ is assumed. Repeating their exercise for a typical RFP indicates $(B_{\text{r}}^{\text{local}}/B_0)_{\text{rms}} \geq 10^{-4}$ would produce stochasticity. The point we make is that even such a level is undetectable, so that Ockham's Razor would favor stochasticity as the cause of observed, nonradiative electron heat loss ($\tau_{Ee} \approx 10^{-4} \text{ s}$) in ZT-40M.

If we assume ZT-40M is stochastic, then the electron heat diffusivity⁹ required to cause τ_{Ee} can be used to estimate the magnetic field-line diffusivity D_F . Krommes et al.,¹⁰ suggest that this estimate will be a lower bound for D_F . If we write $\tau_{Ee} \approx a^2/D_e$, the electron-heat diffusivity (with $a = 0.2 \text{ m}$) is $D_e \approx 4 \times 10^2 \text{ m}^2 \text{ s}^{-1}$. An upper bound¹⁰ on the stochasticity-induced electron-heat diffusivity⁹ is $D_e \approx v_{Te} D_F$. Using $T_e \approx 200 \text{ eV}$ so that $v_{Te} \approx 6 \times 10^6 \text{ ms}^{-1}$, we get $D_F \approx 7 \times 10^{-5} \text{ m}^2 \text{ s}^{-1}$ as a lower bound on the magnetic-field-line diffusivity.

How far does an electron wander during one mean-free-path across the flux surfaces, if indeed $D_F \approx 7 \times 10^{-5} \text{ m}^2 \text{ s}^{-1}$? The most probable electron ($v = v_{Te}/\sqrt{2} \approx v_0$) has a mean-free-path (in a Lorentz plasma with $Z = 1$, $n = 2 \times 10^{19} \text{m}^{-3}$, and $T_e = 200 \text{ eV}$) $\lambda_0 = 20 \text{ m}$. The more relevant number,

though, is λ averaged over j , and this can be shown¹¹ to be $\lambda_j \equiv \int \lambda dj / \int dj = 20 \lambda_0$ for a Lorentz plasma owing to the weighting of suprathermal electrons in carrying j . Using $\lambda_j = 400$ m, we obtain an electron wander $(\Delta x)_j = (2D_F \lambda_j)^{1/2} = 0.3$ m as a lower bound. Thus, in one mean-free-path the j -weighted electron radial wander is similar to the plasma radius!

Consider a slab-geometry RFP with x the normal to "flux surfaces" (like r in a cylinder). The configuration is sustained by a steady, uniform applied E_z . The local magnetic-field-aligned electric field is $E_{||}(x) = E_z B_z(x)/B$. The average gradient length $E_{||}/(\partial E_{||}/\partial x)$ in an RFP will be smaller than a . Thus in ZT-40M, tail electrons wander all over the $E_{||}$ -gradient in one mean-free-path. This voids a local Ohm's law. More importantly, it suggests that RFP sustainment on ZT-40M may be due to export of electron field-aligned momentum from the core (where $E_{||} > j_{||}/\sigma_{||}$) to the outer region (where $E_{||} \leq 0 < j_{||}/\sigma_{||}$).

We have recently developed¹¹ a formal procedure for treating electron-momentum export down the $E_{||}$ -gradient. The treatment is facilitated by some simplifying assumptions (none of which, though, is required for the basic mechanism to be viable):

1. The plasma is isothermal and isodense, and $f^{(0)}(\vec{v})$ is a Maxwellian.
2. Slab-geometry is employed, and $|\vec{B}|$ is uniform.
3. Coulomb scattering is approximated by electron collisions only with massive ions (Lorentz gas).
4. The applied electric field is weak: $E_{||} \ll E_c$, where $E_c \equiv$ critical (runaway) field.¹²
5. $L_F \ll \lambda$ where L_F is the (Kolmogorov) correlation length⁹ and λ is the electron mean-free-path.

In these conditions we have obtained¹¹ the following results:

First: The perturbation $f^{(1)}(\vec{v}, x)$ in the electron distribution function is laminar, depending on x (the normal to "flux surfaces") but not on y or z .

Second: The perturbation $f^{(1)}(\vec{v}, x)$ is purely odd in $\cos\theta$ (where θ is the angle between \vec{v} and \vec{B}); this leads to export of field-aligned momentum, but not of electron number density, down the $E_{||}$ -gradient.

Third: The spatial gradient $\partial f^{(1)}(\vec{v}, x)/\partial x$ causes a Fick's Law flux $-D_e \partial f^{(1)}(\vec{v}, x)/\partial x$, which carries the electron momentum exported down the $E_{||}$ -gradient.

Fourth: For each electron velocity \vec{v} , $f^{(1)}(\vec{v}, x)$ is a solution of a separate Boltzmann equation:

$$f^{(1)}(\vec{v}, x) = - \frac{E_{||}(x)}{E_c} \left(\frac{v}{v_0} \right)^4 \cos\theta f^{(0)}(\vec{v}) + 2\lambda_0 \left(\frac{v}{v_0} \right)^4 |\cos\theta| \frac{\partial}{\partial x} [D_F(x) \frac{\partial f^{(1)}(\vec{v}, x)}{\partial x}]. \quad (1)$$

The first term on the rhs of Eq. (1) is the local Spitzer-Härm¹³ Lorentz-gas solution. The second term on the rhs is (minus) the divergence of the Fick's law flux down the spatial gradient of $f^{(1)}(\vec{v}, x)$. The $(v/v_0)^4 |\cos\theta|$ weighting is caused by the mean-free-path's dependence on \vec{v} .

We solve Eq. (1), with $E_{||}(x)$ and $D_F(x)$ profiles as inputs, at each of 39 velocities (3 angles, θ , at each of 13 speeds, v). The solutions are multiplied by $-\cos\theta$ and integrated $d\vec{v}$ with splines to give $j_{||}(x)$. The contrived boundary condition at the wall is $\partial f^{(1)}/\partial x|_{x=a} = 0$, corresponding to zero momentum export from the plasma to the wall. The $E_{||}(x)$ profile shape is affected by the $j_{||}(x)$ result, because $j_{||}(x)$ controls the magnetic field orientation (via Ampere's law), and $E_{||}(x) = E_z B_z(x)/B$. Thus we iterate the solution of Eq. (1), at each step using an updated $E_{||}(x)$ profile, until the current $j_{||}(x)$ satisfies both $f = ma$ [Eq. (1)] and Ampere's law.

The parameters which we may choose are $\lambda_0 D_F/a^2$ (characterizing the electron wander) and $j_{||}(0)/B$ (corresponding to how hard we push the system). In order to compare with RFP phenomenology we may use $B_y(a)/\langle B_z \rangle$

(corresponding to the pinch parameter, θ) as the second parameter instead of $j_{\parallel}(0)/B$.

A self-consistent solution with uniform diffusivity ($\lambda_0 D_F/a^2 = 0.05$) and pinch parameter $B_y(a)/\langle B_z \rangle = 2.10$ is shown in Fig. 1. The $E_{\parallel}(x)$ profile has the same shape as the $B_z(x)$ profile. Despite the $E_{\parallel}(x)$ profile's sign reversal (at $x = 0.8a$), the field-aligned current $j_{\parallel}(x)$ is almost flat, and never reverses sign. The microscopic reason for this is the spatially diffused profiles of $f^{(1)}(\vec{v}, x)$, shown in Fig. 2. For almost-perpendicular ($\cos\theta = 0.3$) and low-speed ($v/v_0 = 0.8$) velocities, the conduction $f^{(1)}(\vec{v}, x)$ closely resembles $E_{\parallel}(x)$ in shape (Fig. 2, top). However, field-aligned ($\cos\theta = 1.0$) suprathermal electrons ($v/v_0 > 1$) have more diffused $f^{(1)}(\vec{v}, x)$ profiles (Fig. 2, bottom).

In Fig. 3 we show the "resistive anomaly," that is, the ratio of E_z to $j_{\parallel}(0)/\sigma$, where $\sigma \equiv$ nominal local Ohm's-law conductivity. Our resistive anomaly is understated because we do not consider electron-momentum loss to the wall.

An "F- θ diagram" for slab geometry is shown in Fig. 4, using various spatially uniform diffusivities $\lambda_0 D_F/a^2$. The extreme case ($\lambda_0 D_F/a^2 = \infty$) would be called "fully relaxed," and the others "partially relaxed" in dynamo parlance. In our theory of nonlocal conductivity, however, "relaxation" plays no role; instead, the F- θ trajectory is controlled by the range of electron wander, measured by $\lambda_0 D_F/a^2$.

We have also calculated RFP states for tapered profiles of $D_F(x)$, in which D_F is high on axis ($x=0$) but falls to the edge ($x=a$). [This $D_F(x)$ profile may be appropriate to RFP experiments owing to the tendency of the nearby conducting shell to reduce B_z -fluctuations near the edge.] We find that the $j_{\parallel}(x)$ profile responds by also becoming reduced at the edge. This may account for the "Modified" (i.e., tapered at edge) current profiles inferred in experiments.²

Finally, the nonlocal-conductivity model offers some insight on the time scale required for an RFP discharge to relax following a step change in some boundary condition (e.g., toroidal flux or toroidal voltage): Although the model described above is steady-state, it is clear that the $j_{\parallel}(x)$ profile can relax no more quickly than a j -weighted electron-ion collision time.

1. G. L. Siscoe et al., invited paper in session E of AGU Chapman Conf. on Magnetic Reconnection, Los Alamos National Laboratory (1983).
2. D. A. Baker et al., in Plasma Phys. and Contr. Nucl. Fusion Res. 1982, 9th Conf. Proc., Vol. 1, Nuclear Fusion, Suppl. 1983, IAEA, Vienna, p. 587.
3. T. R. Jarboe et al., Phys. Rev. Lett. 51, p. 39 (1983).
4. E. J. Caramana et al., Los Alamos National Laboratory document LA-UR-83-2284 (1983).
5. A. Sykes and J. A. Wesson, in Proc. 8th European Conf. on Contr. Fusion and Plasma Phys., Prague 1977.
6. A. Y. Aydemir and D. C. Barnes, Institute for Fusion Studies Report #102 (1983).
7. J. A. Wesson, private communication 1982.
8. D. C. Barnes, private communication 1983.
9. A. B. Rechester and M. N. Rosenbluth, Phys. Rev. Lett. 40, p. 38 (1978).
10. J. A. Krommes et al., J. Plasma Phys. 30, p. 11 (1983).
11. Abram R. Jacobson and Ronald W. Moses, Los Alamos National Laboratory document LA-UR-83-3440 (1983).
12. H. Knoepfel and D. A. Spong, Nucl. Fusion 19, p. 785 (1979).
13. L. Spitzer and R. Häm, Phys. Rev. 89, p. 977 (1953).




Fig. 1. Normalized profiles of magnetic fields and field-aligned current density for uniform diffusivity.

$$\frac{\lambda_0 D_F}{a^2} = 0.05$$

$$\frac{B_y(a)}{\langle B_z \rangle} = 2.10$$




Fig. 2. Normalized profiles of electron distribution-function perturbation for four velocities, in conditions of Fig. 1.




Fig. 3. Resistive anomaly factor versus pinch parameter, for various diffusivities.




Fig. 4. P- θ trajectories for various diffusivities, in slab-geometry.